

**UNIVERSIDADE ESTADUAL DO MARANHÃO
CENTRO DE CIÊNCIAS AGRÁRIAS
PROGRAMA DE PÓS-GRADUAÇÃO EM AGROECOLOGIA
CURSO DE MESTRADO EM AGROECOLOGIA**

JESÚS ENRIQUE BURGOS GUERRERO

**DIVERSIDADE E COMPOSIÇÃO DA MACROFAUNA DO SOLO NO SISTEMA
AGROFLORESTAL QUESUNGUAL, NA AGRICULTURA DE CORTE QUEIMA E
EM SUCESSÕES FLORESTAIS NA AMAZÔNIA ORIENTAL**

São Luís - MA

2017

JESÚS ENRIQUE BURGOS GUERRERO
Engenheiro Agrônomo

**DIVERSIDADE E COMPOSIÇÃO DA MACROFAUNA DO SOLO NO SISTEMA
AGROFLORESTAL QUESUNGUAL, NA AGRICULTURA DE CORTE QUEIMA E
EM SUCESSÕES FLORESTAIS NA AMAZÔNIA ORIENTAL**

Dissertação apresentada ao Programa de Pós-Graduação em Agroecologia da Universidade Estadual do Maranhão, para a obtenção do título de Mestre em Agroecologia.

Orientador: Prof. Dr. Guillaume Xavier Rousseau

São Luís - MA
2017

Guerrero, Jesús Enrique Burgos.

Diversidade e composição da macrofauna do solo no sistema agroflorestal Quesungual, na agricultura de corte queima e em sucessões florestais na Amazônia Oriental / Jesús Enrique Burgos Guerrero. – São Luís, 2017.

51 f.

Dissertação (Mestrado) – Programa de Pós-Graduação em Agroecologia, Universidade Estadual do Maranhão, 2017.

Orientador: Prof. Dr. Guillaume Xavier Rousseau.

1. Macrofauna. 2. Floresta secundária. 3. Riqueza. 4. Densidade.

I. Título.

CDU 631.95(811.3)

**UNIVERSIDADE ESTADUAL DO MARANHÃO
CENTRO DE CIÊNCIAS AGRÁRIAS
PROGRAMA DE PÓS-GRADUAÇÃO EM AGROECOLOGIA
CURSO DE MESTRADO EM AGROECOLOGIA**

JESÚS ENRIQUE BURGOS GUERRERO

**DIVERSIDADE E COMPOSIÇÃO DA MACROFAUNA DO SOLO NO SISTEMA
AGROFLORESTAL QUESUNGUAL, NA AGRICULTURA DE CORTE QUEIMA E
EM SUCESSÕES FLORESTAIS NA AMAZÔNIA ORIENTAL**

Aprovada em: / /

BANCA EXAMINADORA:

Prof. Dr. Guillaume Xavier Rousseau - UEMA

Profa. Dra. Danielle Camargo Celentano Augusto - UEMA

Prof. Dr. Glécio Machado Siqueira - UFMA

DEDICO

Este trabalho é dedicado para: minha mãe, CLEMENTINA DE JESÚS GUERRERO DIAZ; meu pai, JORGE ENRIQUE BURGOS CABRERA e meu irmão ERNESTO CAMILO BURGOS GUERRERO. Infinita Gratidão pela vida, inspiração, companhia e incondicional apoio. Igualmente gratidão para meus primos e amigos, todos estão no meu coração.

AGRADEÇO

A energia criadora que faz que todas as coincidências aconteçam, e a todas as pessoas lindas e maravilhosas que conheci, nesse país maravilhoso, nesse continente fantástico e nesse mundo único. Quero agradecer especialmente à Guillaume Xavier Rousseau pela paciência, confiança e orientação nesta caminhada. Aos companheiros de caminhada e equipe de trabalho: Luis Hernández, Ernesto Cardozo, Julio Bravo, Henry Reyes, Stefania Pinzon, Sandriel Costa, Jhonatan Muñoz, Robinso Cabezas, Kellen Silva, Thays Frazão, Laiana Leite, Thiago Oliveira. Sem vocês este trabalho não teria sido possível. Aos amigos de tantos momentos especiais: Monica Moreno, Natalia Espitia, Samara Abreu, Gessearia Reis, Livia Reis, Isabelle Victória, Francisneide Loureço, Zélia Nuñez, Raysa Carvalho, Caroline Coelho, Yanna Reis, Ricardo Cueva, Rauricio Barbosa, André Raposo, aos amigos da “Fábula”. Todos vocês e os que me faltam por escrever aqui, já fazem parte da minha família, por que a família não só é de sangue, também de coração. À Danielle Celentano, Rayanne Cristine, Camila Nobre Pinheiro, Marcio Leite e Glecio Machado Siqueira pelas correções, comentários e apoio.

Aos meus colegas da turma de Pós-Graduação em Agroecologia.

Aos habitantes de Alcântara, especialmente aos moradores das comunidades: Pepital, Só Assim, Espera, Cajueiro, Maruda, Mamuna, por sua colaboração e cordialidade com nosso grupo de trabalho. Esta pesquisa é para vocês.

À Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) pela concessão da bolsa de estudos.

À Fundação de Amparo à Pesquisa e ao Desenvolvimento Científico e Tecnológico do Maranhão (FAPEMA) pelo financiamento da pesquisa.

*Eu sei de um país verde como uma maçã
Doce país de milho e laranjas
Sei de um país de palmeiras e montanhas
Doce país que tem sabor de coco e cana
Te levo, te sinto, te amo e te quero*

*E vou te cuidar com tua água e teu ar
E eu vou me educar, porque a terra não se canse
E eu vou te cuidar por tuas florestas e teus mares
E eu vou me educar para te querer sempre e sempre mais.*

*Sei de um país verde como uma maçã
Doce lugar de milho e laranjas
Doce lugar, eu quero vê-lo sempre bem
Sempre alegre, sempre verde, sorrir e florescer*

Ricardo Williams

SUMÁRIO

	Pag
LISTA DE FIGURAS.....	viii
LISTA DE TABELAS.....	ix
RESUMO.....	x
RESUMEN.....	xi
CAPÍTULO I.....	12
1 INTRODUÇÃO.....	13
2 REVISÃO DE LITERATURA	14
2.1 Macrofauna do solo.....	14
2.2 Principais grupos taxonômicos da macrofauna do solo e sua importância como indicadores ambientais.....	14
2.3 Agricultura de corte e queima	16
2.3.1 Efeito do fogo no solo.....	17
2.4 Alternativas ao sistema de agricultura de corte e queima	17
2.5 Sistema Agroflorestal Quesungual.....	18
2.5.1 Vantagens do Sistema Agroflorestal Quesungual.....	19
3 OBJETIVO.....	20
4 HIPÓTESE.....	20
5 REFERÊNCIAS BIBLIOGRÁFICAS.....	21
CAPÍTULO II.....	28
CAPITULO II: Quesungual Agroforestry System conserves soil macrofauna density and richness of the Amazon forest.....	29
ABSTRACT.....	29
INTRODUCTION.....	30
MATERIALS AND METHODS.....	31
RESULTS.....	36
DISCUSSION.....	42
CONCLUSIONS.....	44
ACKNOWLEDGMENTS.....	44
REFERENCES.....	45
SUPPLEMENTARY MATERIAL.....	50

LISTA DE FIGURAS

	Pag
Figura 1 Location of study areas in Alcântara County, Maranhão state, Brazil.....	32
Figura 2 The eight land-use evaluated in Alcântara County, Maranhão state, Brazil.....	34
Figura 3 Density of soil macrofauna (ind/m ²) by taxa and dendogram of Bray-Curtis index in Alcântara County, Maranhão state, Brazil.....	39
Figura 4 Canonical redundancy analysis (RDA) with soil macrofauna density, soil physico-chemical and sampling sites in different land-uses in Alcântara County, Maranhão state, Brazil	41
Figura 5 Taxa accumulation curve of soil macrofauna under different land uses in Alcântara County, Eastern Amazon.....	51

LISTA DE TABELAS

	Pag
Tabela 1 Description of plots and land uses in Alcântara County, Maranhão state, Brazil.....	33
Tabela 2 Soil macrofauna community-structure indices (mean and standard errors) in different land uses within Alcântara County, Maranhão state, Brazil	37
Tabela 3 Physical and chemical attributes of soils (average and standard errors) in different land use in Alcântara County, Maranhão state, Brazil	40
Tabela 4 Means of density (Ind/m ²) of the soil macrofauna by taxa and functional groups (average and standard errors) under different land uses in Alcântara County, Eastern Amazon....	50

RESUMO

O Sistema Agroflorestal Quesungual (QAS) originário da América Central, pode ser uma alternativa sustentável para os produtores familiares de pequena escala na Amazônia Oriental, onde a intensificação da agricultura com corte e queima ameaça o meio ambiente e sua subsistência. Este é o primeiro experimento para testar esse sistema na região amazônica. Para a avaliação do seu impacto na qualidade do solo nos determinamos a abundância e riqueza da macrofauna do solo e os atributos físicos e químicos do solo no QAS, agricultura de corte e queima (SB) e capoeiras (SF) de diferentes idades de pousio em comparação com florestas maduras (MF) no município de Alcântara, estado do Maranhão. Em total 32 áreas (40 X 50 m) foram avaliadas; as análises de dados incluíram modelos mistos, cluster e RDA. As comunidades da macrofauna foram sensíveis indicadores do sistema do uso do solo, especialmente predadores, organismos saprófagos e transformadores da serapilheira. Os resultados indicam que QAS e MF conservam similar densidade e riqueza da macrofauna, enquanto SB tem um forte impacto negativo sobre ela. O Sistema Agroflorestal Quesungual maximiza a acumulação de carbono orgânico no solo e representa uma alternativa viável para a promoção da conservação da biodiversidade e da segurança alimentar na Amazônia Oriental.

Palavras chave: Macrofauna, floresta secundária, riqueza, densidade, Amazônia.

RESUMEN

El Sistema Agroforestal Quesungual (QAS) originario de América Central puede ser una alternativa sustentable para agricultores familiares de pequeña escala en la Amazonia Oriental, donde la intensificación de la agricultura con corte y quema amenaza el medio ambiente y su subsistencia. Este es el primer experimento para probar este sistema en la región Amazónica. Para evaluar su impacto en la calidad del suelo determinamos la abundancia y diversidad de la macrofauna, así como los atributos físicos y químicos del suelo en el QAS, agricultura con corte y quema (SB) y bosques secundarios (SF) de diferentes edades de barbecho en comparación con bosques maduros en el municipio de Alcântara, estado de Maranhão. En total 32 áreas (40 X 50 m) fueron evaluadas. El análisis de los datos incluyó modelos mixtos, clusters y RDA. La macrofauna fue un sensible indicador del sistema de uso de la tierra, especialmente predadores, organismos saprófagos y transformadores de la hojarasca. Los resultados indican que QAS y MF conservan similar riqueza y diversidad de la macrofauna, mientras que SB tiene un fuerte impacto negativo en ella. El Sistema Agroforestal Quesungual maximiza la acumulación de carbono orgánico en el suelo y representa una alternativa viable para promover la conservación de la biodiversidad del suelo y la seguridad alimentaria en la Amazonia Oriental.

Palabras clave: Macrofauna, bosque secundario, riqueza, densidad, Amazonia.

CAPÍTULO I

1 INTRODUÇÃO

Na atualidade são muitos os esforços para substituir o uso do sistema de agricultura de corte e queima (PEDROSO et al., 2008). Esse sistema é sustentável e até pode favorecer a biodiversidade (PEDROSO et al., 2008; PADOCH; PINEDO-VASQUEZ, 2010; LAWRENCE et al., 2010), mas para seu adequado funcionamento precisa de alta disponibilidade de terras e prolongados períodos de pousio. (BARRIOS et al., 2005; PEDROSO et al., 2008; PADOCH; PINEDO-VASQUEZ, 2010; LAWRENCE et al., 2010), condições cada vez mais difíceis de conseguir no atual contexto de forte pressão demográfica (METZGER, 2003; MERTZ et al., 2009; VAN VLIET et al., 2012).

Para sua substituição várias alternativas têm sido propostas (PEDROSO et al., 2008), muitas delas baseadas no uso de sistemas agroflorestais. Uma dessas alternativas é o chamado Sistema Agroflorestal Quesungual, desenvolvido e praticado em Honduras, América Central, por comunidades indígenas há séculos (HELLIN et al., 1999). O sistema foi criado para a gestão do solo em áreas de encosta, com solos facilmente degradáveis, períodos de seca prolongados e onde o fogo era tradicionalmente utilizado para a limpeza das áreas de cultivo.

Embora na Amazônia Oriental, não sejam encontrados declives acentuados ou encostas, todos os outros fatores que caracterizam o Sistema Agroflorestal Quesungual estão presentes na região de estudo, o que sugere que esse sistema também pode ser viável para essa região da Amazônia.

Por outro lado a densidade e diversidade da macrofauna do solo é uma forma eficaz e rápida de avaliar o impacto de novas estratégias de manejo da terra (LOBRY DE BRUYN, 1999) já que estes organismos são fortemente afetados pelas mudanças nas condições do seu ambiente (DECAËNS, 2010; POSTMA BLAAUW et al., 2012; BARETTA et al., 2014).

Nesse contexto, esse trabalho teve como objetivo caracterizar a densidade e a diversidade da macrofauna edáfica no Sistema Agroflorestal Quesungual, em sistema tradicional de cultivo com fogo e nas florestas secundárias e maduras da

Amazônia oriental.

2 REVISÃO DE LITERATURA

2.1 Macrofauna do solo

A macrofauna do solo refere-se aos invertebrados do solo, cujo diâmetro corporal varia entre 2 mm e 20 mm (facilmente visíveis ao olho nú) e que vivem dentro ou sobre a superfície do solo em todas ou pelo menos uma das fases do seu ciclo de vida (KEVAN, 1968; SWIFT et al, 1979; BRAUN et al., 2009). Para seu melhor estudo, esses invertebrados edáficos são divididos em diferentes níveis de identificação taxonômica ou táxons (classes, ordens e famílias) (BRAUN et al., 2009).

Estes organismos são muito importantes já que desempenham um papel chave no funcionamento do solo e dos ecossistemas em geral, modificando seu ambiente através da participação ativa nos ciclos biogeoquímicos, contribuindo no desenvolvimento estrutural do solo (SILVA et al., 2006) e aumentando a porosidade, agregação e retenção da água, através da formação de estruturas como túneis ou câmaras (LAVELLE; SPAIN, 2001; AIRA et al., 2008). Por isso, em alguns casos são chamados os engenheiros do ecossistema (LAVELLE et al., 2006).

A presença das populações da macrofauna do solo nos agroecossistemas é um fator importante para a sustentabilidade dos sistemas agrícolas (BRUSSAARD et al., 2007), já que esses organismos além de serem atores nas funções do solo (MARICHAL et al, 2014) também podem servir como indicadores das mudanças no seu ambiente (VASCONCELLOS et al., 2013; MARICHAL et al., 2014).

2.2 Principais grupos taxonômicos da macrofauna do solo.

Alguns dos grupos taxonômicos da macrofauna do solo mais importantes por sua alta abundância e sensibilidade aos distúrbios ou mudanças no ambiente em que vivem são:

O ordem Isóptera (cupins) e a família Formicidae (formigas), são insetos sociais, amplamente distribuídos no mundo e com alta capacidade de adaptação a diferentes tipos de habitats. Mas também são sensíveis a prolongadas ou extremas alterações do ambiente, variando em sua abundância, dominância ou diversidade (BAUTISTA et al., 2009). São utilizados em estudos de biodiversidade, monitoramento, fragmentação e processos de restauração (BACCARO, 2006; COSTA et al., 2100; PAIS; VARANDA, 2010; ROUSSEAU et al., 2010).

A Classe Oligochaeta (minhocas) é fundamental no funcionamento do solo por sua estreita relação com a dinâmica da matéria orgânica (FRAGOSO et al., 1997) e por seu aporte direto na estrutura do solo com sua movimentação no perfil, e a formação de estruturas biogênicas altamente estáveis (BLANCHART et al., 2004). Sendo altamente influenciadas por práticas de manejo do solo (BARTZ et al., 2014) e favorecidas pela presença de matéria orgânica e cobertura morta (FONTE et al., 2009).

O ordem Coleóptera (besouros) é um dos mais diversos grupos no total de insetos descritos no mundo (ORGIAZZI et al., 2016). Eles contribuem diretamente nos processos de decomposição da matéria orgânica, ciclagem de nutrientes e movimentação do solo, sendo capazes de explorar uma alta diversidade de nichos ecológicos (ORGIAZZI et al., 2016).

Por sua vez, predadores como Araneae (aranhas), Opilionida, Chilopoda (piochos de cobra) e Pseudoscorpionida são especialmente relacionados com ecossistemas estáveis e a presença desses grupos é sinal de um fluxo normal de energia dentro da cadeia trófica do ecossistema. (ROUSSEAU et al., 2014)

Outros invertebrados como Isopodos (tatuzinhos), Diplopodos (centopéias), Gastropodos (lesmas e caracóis) também são amplamente sugeridos como indicadores do funcionamento do solo, por sua estreita relação com ambientais mais complexos e estáveis, com maior conteúdo de carbono orgânico e pela sensibilidade aos distúrbios naturais ou antropogênicos (VASCONCELLOS et al., 2013; MORON-RIOS et al., 2010; BARETTA et al., 2011; TIZADO; NUÑES-PEREZ, 2016).

2.3 Agricultura de corte e queima

Há milênios na região amazônica, a conversão das áreas florestais primárias ou secundárias em áreas de cultivos agrícolas se faz através da chamada agricultura com corte e queima (PEDROSO et al., 2008; BARNI et al., 2012). Esse é um sistema sustentável quando é praticado da forma correta, em grandes áreas florestais, com baixas densidades populacionais, adequados sistemas de plantação e longos períodos de pousio (KLEINMAN et al., 1995; JOHNSON et al., 2001; MENDOZA-VEGA et al., 2003; PEDROSO et al., 2008). Sendo capaz de manter ou inclusive promover a biodiversidade (PEDROSO et al., 2008).

Seus componentes básicos são o corte da vegetação primária ou secundária, a queima do material vegetal cortado, o cultivo e o pousio ou período de regeneração da vegetação cortada (KLEINMAN et al., 1995; PEDROSO et al., 2008). O período de cultivo geralmente tem uma duração de um até dois anos, logo, o período de pousio tem que ser muito mais longo que o período de cultivo (KLEINMAN et al., 1995; PEDROSO et al., 2008; MERTZ et al., 2009).

Entre as vantagens dessa prática está a facilitação dos tratamentos culturais após a queimada, a eliminação de ervas espontâneas, pragas, doenças e principalmente a adição de nutrientes constituintes da cinza que melhoram as propriedades químicas do solo (KLEINMAN et al., 1995; PEDROSO et al., 2008; MERTZ et al., 2009).

No entanto, no atual contexto de alta pressão demográfica e alta demanda de terras (MERTZ et al., 2009; VAN VLIET et al., 2012), os adequados períodos de pousio são cada vez menos praticados (COOMES et al., 2000; METZGER 2003; LAWRENCE et al., 2010), com importantes consequências na sustentabilidade do sistema. Além disso a agricultura com corte e queima é questionada por sua participação no processo de desmatamento, perda de biodiversidade (PEDROSO et al., 2008), aquecimento global devido as emissões de CO₂ e degradação dos solos. (COE et al., 2013),

2.3.1 Efeito do fogo no solo

No que se refere ao solo, o uso do fogo tem importantes consequências nas propriedades físicas, químicas e biológicas (MOURA et al., 2009; LAWRENCE et al., 2010; ROSSI et al., 2010). Principalmente pela alta temperatura que pode alcançar o solo (400 °C) (MATLACK, 1993; GIMENO-GARCÍA et al., 2004), evaporação da água e a eliminação da matéria orgânica (VERMA; JAYAKUMAR, 2012).

O pH pode aumentar pela adição das cinzas, caracterizadas por ter alto pH (MOLINA et al., 2007). A densidade aparente aumenta devido ao colapso dos agregados, afetando negativamente a porosidade e permeabilidade do solo (CERTINI, 2005). Os componentes da textura do solo (areia, silte e argila) têm limiares de alta temperatura e geralmente não são afetados pelo fogo (VERMA; JAYAKUMAR, 2012),

O conteúdo de nutrientes do solo pode variar segundo o tipo de elemento, mas geralmente resulta em um incremento na sua disponibilidade (KUTIEL; NAVEH, 1987). O carbono orgânico diminui, o cálcio aumenta e os estoques de magnésio, potássio e fósforo permanecem sem maiores variações (GARCÍA-OLIVA et al., 1999; NEFF et al., 2005; MERTZ et al., 2009).

Por sua vez, a biota do solo diminui consideravelmente. Principalmente as populações de organismos como fungos e bactérias (DAHLBERG et al., 2001) e comunidades de invertebrados (BARROS et al., 2002; BARRIOS et al., 2005; SAINT-GERMAIN et al., 2005; BUDDLE et al., 2006), os quais sucumbem diretamente pelas chamas e gases quentes ou indiretamente pela destruição de suas fontes de comida e abrigo (MATHIEU et al., 2005; ROSSI et al., 2010).

2.4 Alternativas ao sistema de agricultura de corte e queima.

Na atualidade muitos governos e organizações no mundo inteiro fazem esforços para a substituição da agricultura de corte e queima por práticas menos agressivas ao meio ambiente (PEDROSO et al., 2008; COE et al., 2013). Várias alternativas tem sido sugeridas, entre elas o uso de coberturas mortas

(NORGROVE; HAUSER, 2002; VIELHAUER et al., 2004; DENICH et al., 2005), o cultivo em aléias (MOURA et al 2014), o plantio direto (BARTZ et al., 2014) e a implementação de Sistemas Agroflorestais ou SAFs (DASH; MISRA, 2001; FAGERSTRÖM et al., 2002, VIELHAUER et al., 2004; PINHO et al., 2012).

Os SAFs tem especial destaque por serem um sistema de produção que favorece a diversidade de plantas, a conservação do solo, a provisão de serviços ambientais e que dá considerável renda para os agricultores (WELCHES et al., 2008; ROUSSEAU et al., 2013; CARDOZO et al., 2015).

O Centro Mundial Agroflorestal (ICRAF), define os SAFs como sistemas e tecnologias de uso da terra que combinam árvores florestais com culturas agrícolas, pastagens e ou animais, dependendo do tempo e espaço para aumentar e otimizar a produção de forma constante (NAIR, 1993).

2.5 Sistema Agroflorestal Quesungual

Entre as alternativas foi proposto um sistema praticado por comunidades indígenas de Honduras, América Central, há centenas de anos (HELLIN et al., 1999). Esta antiga prática agroflorestal foi observada e descrita por pesquisadores da FAO na década de 1990 e chamada sistema agroflorestal Quesungual pela comunidade onde foi identificado pela primeira vez (HELLIN et al., 1999; PAULI et al., 2011).

Este é um sistema de produção para pequenos agricultores, que inclui um pacote de tecnologias para o manejo sustentável da vegetação, do solo e da água em áreas de encosta com tendência de seca no trópico sub-úmido (CIAT, 2009).

Esse sistema sustenta-se em quatro princípios básicos (CIAT, 2010)

- Não corte e queima, mediante a gestão corte e poda (parcial, seletiva e progressiva) da vegetação nativa.
- Cobertura permanente do solo, por meio da deposição contínua de biomassa

de árvores, arbustos, ervas espontâneas e ou resíduos de culturas.

- Mínima perturbação do solo através do plantio direto.
- Uso eficiente de fertilizantes, através da sua aplicação adequada no tempo, tipo, quantidade e localização.

Para sua implementação, as árvores e arbustos com algum tipo de interesse madeireiro ou frutíferos são deixados em pé, enquanto que, o material vegetal restante é cortado a uma altura de 1,5 metros, sem machucar o tronco nem a raiz, para permitir sua regeneração posterior. Todo o material podado é cortado e usado como cobertura morta sobre a área total de cultivo (FAO, 2005).

2.5.1 Vantagens da implementação do Sistema Agroflorestal Quesungual

Existem diversas vantagens na implementação deste sistema (CIAT, 2010).

- Incremento paulatino da produtividade, garantindo a segurança alimentar dos camponeses.
- Constante fornecimento de lenha para uso doméstico.
- Melhoria na prestação de serviços ecossistêmicos por incremento no sequestro de carbono e diminuição da erosão do solo, redução do desmatamento e da emissão de gases efeito estufa;
- E a conservação da biodiversidade local e a proteção do recurso água, solo e vegetação.

Além disso, vários estudos afirmam que esse sistema permite aproveitar a área de cultivo por mais tempo (> 7 anos) enquanto que nos sistemas de agricultura com corte e queima após 2 ou 3 anos as áreas são abandonadas (PAULI et al., 2011).

3. OBJETIVO

Determinar e comparar a densidade e diversidade da macrofauna do solo no Sistema Agroflorestal Quesungual, na agricultura com corte-queima, em florestas secundárias e em florestas maduras da Amazonia oriental.

4. HIPÓTESE

O Sistema Agroflorestal Quesungual pode manter a mesma densidade e diversidade da macrofauna do solo que as áreas florestais melhor conservadas.

5. REFERÊNCIAS BIBLIOGRÁFICAS

AIRA, M.; SAMPEDRO, L.; MONROY, F.; DOMÍNGUEZ, J. Detritivorous earthworms directly modify the structure, thus altering the functioning of a microdecomposer food web. **Soil Biology and Biochemistry**, v. 40, n. 10, p. 2511-2516, 2008.

BACCARO, F. B. Hymenoptera: Formicidae. Instituto Nacional de Pesquisas da Amazônia – INPA. Programa de Pesquisa em Biodiversidade - PPBIO. Faculdades Cathedral. 36 p, 2006.

BARETTA, D.; BARTZ, M.; FACHINI, I.; ANSELMINI, R.; ZORTÉA, T.; BARETTA, C. Soil fauna and its relation with environmental variables in soil management systems. **Revista Ciência Agronômica**. v. 45, n. 5, p. 871-879, 2014.

BARETTA, D.; SANTOS, J.C.P.; SEGAT, J.C.; GEREMIA, E.V.; DE OLIVEIRA FILHO, L.C.I.; ALVES, M.V. Fauna edáfica e qualidade do solo, in: Klauberg Filho, o.; Mafra, A.L.; Gatiboni, L.C. Tópicos especiais em ciência do solo, Sociedade Brasileira de Ciência do Solo, Viçosa, pp. 141-192, 2011.

BARNI, P. E.; FEARNSIDE, P. M.; GRAÇA, P. M. L. A. Desmatamento no sul do Estado de Roraima: padrões de distribuição em função de Projetos de Assentamento do INCRA e da distância das principais rodovias (BR-174 e BR-210). **Acta Amazônica**, v. 42, n. 2, p. 195-204, 2012.

BARRIOS, E.; COBO, J.G.; RAO, I.M.; THOMAS, R.J.; AMEZQUITA, E.; JIMENEZ, J.J.; RONDON, M.A. Fallow management for soil fertility recovery in tropical Andean agroecosystems in Colombia. **Agriculture Ecosystems and Environment**, v. 110, p. 29–42, 2005.

BARROS, E.; PASHANASI, B.; CONSTANTINO, R.; LAVELLE, P. Effects of land-use system on the soil macrofauna in western Brazilian Amazonia. **Biology and Fertility of Soils**, v. 35, p. 338–347, 2002.

BARTZ, C.; BROWN, G.; DA ROSA, M.; FILHO, O.; JAMES, S.; DECANS, T.; BARETTA, D. Earthworm richness in land use systems in Santa Catarina, Brazil. **Applied soil ecology** v. 83, p. 59-70, 2014.

BAUTISTA, F.; DIAZ-CASTELAZO, C.; GARCIA-ROBLES, M. changes in soil macrofauna in agroecosystems derived from low deciduous tropical forest on leptosols from karstic zones. **Tropical and Subtropical Agroecosystems**, v. 10, p. 185 – 197, 2009.

BLANCHART, E.; ALBRECHT, A.; BROWN, G. G.; DECÄENS, T.; DUBOISSET, A.; LAVELLE, P.; MARIANI, L.; ROOSE, E. Effects of tropical endogeic earthworms on soil erosion: a review. **Agriculture, Ecosystems and Environment**, v. 104, p. 303-315, 2004.

BROWN, G.; AQUINO, MASCHIO, W.; FROUFE, L. Macrofauna do solo em sistemas agroflorestais e mata atlântica em regeneração nos municípios de Barra do Turvo, SP, e Adrianópolis, PR. Colombo, Embrapa Florestas. 51 p. 2009.

BRUSSAARD, L.; DE RUITER, P.C.; BROWN, G. Soil biodiversity for agricultural sustainability. **Agriculture Ecosystem and Environment**. v. 121, p. 233–244, 2007.

BUDDLE, C.M.; LANGOR, D.W.; POHL, G.R.; SPENCE, J.R. Arthropod responses to harvesting and wildfire: Implications for emulation of natural disturbances in forest management. **Biological Conservation** v. 128, p. 346-357, 2006.

CARDOZO, E.; MUCHAVISOY, H.; SILVA, H.; ZELARAYAN, M.; LEITE, M.; ROUSSEAU, G. X.; GEHRING, C. Species richness increases income in agroforestry systems of eastern Amazonia. **Agroforestry systems**. v. 89, n. 5, p. 901-916, 2015.

CERTINI, G. Effect of fire on properties of soil - A review. **Oecologia**. v. 143, p. 1-10, 2005.

CIAT - CENTRO INTERNACIONAL DE AGRICULTURA TROPICAL. Quesungual slash and mulch agroforestry system (QSMAS): Improving crop water productivity, food security and resource quality in the subhumid tropics. CPWF Project Report. Cali, Colombia. 64 p. 2009.

CIAT - CENTRO INTERNACIONAL DE AGRICULTURA TROPICAL. Quesungual Slash & Mulch Agroforestry System: An Eco-Efficient Option for the Rural Poor. 2010.

COE, M.; MARTHEWS, T.; COSTA, M.; GALBRAITH, D.; GREENGLASS, N.; IMBUZEIRO, H.; LEVINE, N.; MALHI, Y.; MOORCROFT, P.; MUZA, M.; POWELL, T.; SALESKA, S.; SOLORZANO, L.; WANG, J. Deforestation and climate feedbacks threaten the ecological integrity of south –southeastern Amazonia. **The Royal Society**. 2013.

COOMES, O.; GRIMARD, F.; BURT, G. Tropical forests and shifting cultivation: secondary forest fallow dynamics among traditional farmers of the Peruvian Amazon. **Ecological Economics**, v. 32, p. 109-124, 2000.

COSTA, C.; RIBEIRO, S.; CASTRO, P. Ants as Bioindicators of Natural Succession in Savanna and Riparian Vegetation Impacted by Dredging in the Jequitinhonha River Basin, Brazil. **Restoration Ecology** Vol. 18, n. S1, p. 148–157, 2010.

DAHLBERG, A.; SCHIMMEL, J.; TAYLOR, A.F.S.; JOHANNESSON, H. Post-fire legacy of ectomycorrhizal fungal communities in the Swedish boreal forest in relation to fire severity and logging intensity. **Biological Conservation**, v. 100, p. 151-161, 2001.

DASH, S. S.; MISRA, M. K. Studies on hill agro-ecosystems of three tribal villages on the Eastern Ghats of Orissa, India. **Agriculture, Ecosystems and Environment**, v. 86, n. 1, p. 287-302, 2001.

DECAËNS, T. Macroecological patterns in soil communities. **Global ecology Biogeography**. v. 19, 287-302, 2010.

DENICH, M.; VLEK, P. L. G.; SÁ, T. D.; VIELHAUER, K.; LÜCKE, W. A concept for

the development of fire-free fallow management in the Eastern Amazon, Brazil. **Agriculture, Ecosystems and Environment**, v. 110, p. 43-58, 2005.

FAGERSTRÖM, M. H. H.; NILSSON, S. I.; VAN NOORDWIJK, M.; PHIEN, T.; OLSSON, M.; HANSSON, A.; SVENSSON, C. Does Tephrosia candida as fallow species, hedgerow or mulch improve nutrient cycling and prevent nutrient losses by erosion on slopes in northern Vietnam? **Agriculture, Ecosystems and Environment**, v. 90, p. 291-304, 2002.

FONTE, S.; WINSOME, T.; SIX, J. Earthworm populations in relation to soil organic matter dynamics and management in California tomato cropping systems. **Applied Soil Ecology** 41, 206-214. 2009.

FAO - FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS. El sistema agroforestal Quesungual: Una opción para el manejo de suelos en zonas secas de ladera. 50 p. 2005.

FRAGOSO, C.; BROWN, G.G.; PATRÓN, J.C.; BLANCHART E.; LAVELLE, E.; PASHANASI, B.; SENAPATI, T. Kumar Agricultural intensification, soil biodiversity and agroecosystem function in the tropics " the role of earthworms, **Applied Soil Ecology**, v. 6, p. 17–35, 1997.

GARCIA-OLIVA, F.; SANFORD, JR.; R.L.; KELLY, E. Effects of slash-and-burn management on soil aggregate organic C and N in a tropical deciduous forest. **Geoderma**. v. 88, p. 1–12, 1999.

GIMENO-GARCIA, E.; ANDREU, V.; RUBIO, J. Spatial patterns of soil temperatures during experimental fires. **Geoderma**, v. 118, p. 17–38, 2004.

HELLIN, J.; WELCHEZ, L.A.; CHERRETT, I. The Quezungal System: an indigenous agroforestry system from western Honduras. **Agroforestry Systems**, v. 46, p. 229-237, 1999.

JOHNSON, C. M.; VIEIRA, I. C. G.; ZARIN, D. J.; FRIZANO, J.; JOHNSON, A. H. Carbon and nutrient storage in primary and secondary forests in eastern Amazonia. **Forest Ecology and Management**, v. 147, n. 2, p. 245-252, 2001.

KEVAN, D.K.McE. Soil animals. Northumberland Press, Gateshead on Tyne, UK. 1968.

KLEINMAN, P. J.; PIMENTEL, D.; BRYANT, R. B. The ecological sustainability of slash-and-burn agriculture. **Agriculture, Ecosystems and Environment**. v. 52, n. 2-3, p. 235-249, 1995.

KUTIEL, P.; NAVEH Z. The effect of fire on nutrients in a pine forest soil. **Plant and Soil**. v. 104, p. 269-274, 1987.

LAVALLE, P.; A.V. SPAIN. Soil ecology. Norwell MA: Kluwer Academic Publishers. Dordrecht, The Netherlands, 654 p, 2001.

LAVELLE, P.; DECAËNS, T.; AUBERT, M.; BAROT, S.; BLOUIN, M.; BUREAU, F.; MARGERIE, P.; MORA, P.; ROSSI, J.P. Soil invertebrates and ecosystem services. *European Journal of Soil Biology*, v. 42, p. 3-15, 2006.

LAWRENCE, D.; C. RADEL.; K. TULLY.; B. SCHMOOK.; L. SCHNEIDER. Untangling a decline in tropical forest resilience: constraints on the sustainability of shifting cultivation across the globe. *Biotropica*, v. 42, p. 21–30, 2010.

LOBRY DE BRUYN, L.A. Ants as bioindicators of soil function in rural environments, *Agriculture, Ecosystems and Environment*, v. 74, p. 411–425, 1999.

MARICHAL, R.; GRIMALDI, M.; FEIJOO, A.; OSZWALD, J.; PRAXEDES, C.; RUIZ-COBO, D.; HURTADO, M.; DESJARDINS, H.; DA SILVA, M.; COSTA, G.; MIRANDA, I.; OLIVEIRA, M.; BROWN, J.; TSÉLOUIKO, S.; MARTINS, M.; DECAËNS, T.; VELASQUEZ, E.; LAVELLE, P. Soil macro invertebrate communities and ecosystem services in deforested landscapes of Amazonia. *Applied Soil Ecology*, v. 83, 177–185 p. 2014.

MATHIEU, J.; ROSSI, J.-P.; MORA, P.; LAVELLE, P.; MARTINS, P.F.D.S.; ROULAND, C.; GRIMALDI, M. Recovery of soil macrofauna communities after forest clearance in Eastern Amazonia. Brazil. *Conservation Biology*. v. 19, p. 1598–1605, 2005.

MATLACK, G.R. Microenvironment variation within and among forest edge sites in the eastern United States. *Conservation Biology*. v. 66, p. 185–194, 1993.

MENDOZA-VEGA, J.; KARLTUN, E.; OLSSON, M. Estimations of amounts of soil organic and fine root carbon in land use and land cover classes, and soil types of Chiapas highlands, Mexico. *Forest Ecology and Management*, v. 177, p. 1-16, 2003.

MERTZ, O.; PADOCH, C.; FOX, J.; CRAMB, R.A.; LEISZ, S.J.; LAM, N.I.; VIEN, T.D. Swidden change in Southeast Asia: Understanding Causes and Consequences. *Human Ecology*, v. 37, p. 259-264, 2009.

METZGER, J. P. effects of slash and burn fallow periods on landscapes structure. *Environmental conservation*, v. 30 n. 4, p. 325-333. 2003.

MOLINA, M.; FUENTES, R.; CALDERÓN, R. Impact of forest fire ash on surface charge characteristics of Andisols. *Soil Science*. v. 172, n. 10, p. 820-834, 2007.

MORÓN-RIOS, A; RODRIGUEZ, M.A; PÉREZ-CAMACHO, L.; REBOLLO, S Effects of seasonal grazing and precipitation regime on the soil macroinvertebrates of a Mediterranean old-field, *European Journal Soil Biology*, v. 46. 91-96. 2010.

MOURA, E.G. D.E.; N.G. MOURA, E.S.; MARQUES, K.M.; PINHEIRO, J.R.S. COSTA SOBRINHO.; A.C.F. AGUIAR. Evaluating chemical and physical quality indicators for a structurally fragile tropical soil. *Soil Use and Management*, v. 25 n. 4, p. 368–375, 2009.

MOURA, E.G.; AGUIAR, A.C.F.; PIEDADE, A.R.; ROUSSEAU, G.X. Contribution of legume tree residues and macrofauna to the improvement of abiotic soil properties in the easter Amazon. **Applied Soil Ecology**. v. 86, p. 91-99, 2014.

NAIR, P.K.R. An introduction to Agroforestry. Kluwer Academic Publishers. 499 p. 1993.

NEFF, J.; HARDEN, J.; GLEIXNER, G. Fire effects on soil organic matter content, composition, and nutrients in boreal interior Alaska. **Canadian Journal of Forest Research**. v. 35, p. 2178-2187, 2005.

NORGROVE, L.; HAUSER, S. Yield of plantain grown under different tree densities and 'slash and mulch' versus 'slash and burn' management in an agrisilvicultural system in southern Cameroon. **Field Crops Research**, v. 78, n. 2, p. 185-195, 2002.

ORGIAZZI, A.; BARDGETT, R.D.; BARRIOS, E.; BEHAN-PELLETIER, V.; BRIONES, M.J.I.; CHOTTE, J.-L.; DE DEYN, G.B.; EGGLETON, P.; FIERER, N.; FRASER, T.; HEDLUND, K.; JEFFERY, S.; JOHNSON, N.C.; JONES, A.; KANDELER, E.; KANEKO, N.; LAVELLE, P.; LEMANCEAU, P.; MIKO, L.; MONTANARELLA, L.; MOREIRA, F.M.S.; RAMIREZ, K.S.; SCHEU, S.; SINGH, B.K.; SIX, J.; VAN DER PUTTEN, W.H.; WALL, D.H. Global Soil Biodiversity Atlas. European Commission, Publications Office of the European Union, Luxembourg. 176 pp. 2016.

PADOCH, C.; PINEDO-VASQUEZ, M. Saving slash and burn to save biodiversity. **Biotropica**. v. 42, n. 5, p. 550-552, 2010.

PAIS, M.P.; VARANDA, E.M. Arthropod recolonization in the restauration of a semideciduous forest in southeastern Brazil. **Neotropical Entomology**, v. 39, p. 198-206, 2010.

PAULI, N.; BARRIOS, E.; CONACHER, A.J.; OBERTHÜR, T. Soil macrofauna in agricultural landscapes dominated by the Quesungual slash-and-mulch agroforestry system, Western Honduras. **Applied Soil Ecology**. v. 47, p. 119–132, 2011.

PEDROSO J.; MURRIETA, R.; ADAMS, C. A agricultura de corte e queima: um sistema em transformação. Boletim Museu do Para Emílio Goeldi, Ciências Humanas, Belém, v. 3, n. 2, p. 153-174, 2008.

PINHO R. C.; MILLER R. P.; ALFAIA S. S. Agroforestry and the improvement of soil fertility: a view from Amazonia. **Applaid Environment Soil Science**. 2012.

POSTMA BLAAUW, M.B; GOEDE, R.G.M; BLOEM, J; FABER, J.H; BRUSSARD, L. Agricultural intensification and de-intensification difentially affect taxonomic diversity of predatory mites, earthworms, enchytracids, nematodes and bactéria. **Applaid Soil Ecology**. v. 57, p. 39-49, 2012.

ROSSI, J.P.; CELINI, L.; MORA, P.; MATHIEU J.; LAPIED, E.; NAHMANI, J.; PONGE, J.; LAVELLE, P. Decreasing fallow duration in tropical slash-and-burn agriculture alters soil macroinvertebrate diversity: A case study in southern French GuianaAgriculture, **Agriculture, Ecosystems and Environment**, v. 135, p. 148–154, 2010.

ROUSSEAU, G.X.; P.R. DOS S. SILVA.; C.J.R. DE CARVALHO. Earthworms, ants and other arthropods as soil health indicators in traditional and nofire agroecosystems From Eastern Brazilian Amazonia. **Acta Zoológica Mexicana** (n.s.) (2): 117-134, 2010.

ROUSSEAU G.X.; P.R. DOS S. SILVA.; C.J.R. DE CARVALHO. Impacto do histórico de uso na composição da macrofauna do solo em uma cronoseqüência de capoeiras, florestas e pastos no Centro de Endemismo Belém, Amazônia Oriental. **Acta Amazônica**. v. 44 n. 4, 499-512, 2014.

ROUSSEAU, L.; FONTE, S.; VAN DER HOEK, R.; LAVELLE, P. Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. **Ecological Indicators**. v. 27, p. 71–82, 2013.

SAINT-GERMAIN, M.; LARRIVÉE, M.; DRAPEAU, P.; FAHRIG, L.; BUDDLE, C.M. Short.term response of ground beetles (Coleoptera: Carabidae) to fire and logging in a spruce-dominated boreal landscape. **Forest Ecology and Management**, v. 212, p. 118-126. 2005.

SILVA, R. F.; AQUINO, A. M.; MERCANTE, F. M.; GUIMARÃES, M. Populações de oligoquetos (Annelida: Oligochaeta) em um latossolo vermelho submetido a sistemas de uso do solo. **Ciência Rural**, Santa Maria, v. 36, n. 2, 2006.

SWIFT, M.J.; W. HEAL.; J.M. ANDERSON. Decomposition in terrestrial ecosystems. University of California Press, Berkeley, USA. 1979.

TIZADO, E.; NUÑES-PEREZ, E. Terrestrial arthropods in the initial restoration stages of anthracite coal mine spoil heaps in northwestern Spain potential usefulness of higher taxa as restoration indicators. **Land degrad develop**, v. 21, p. 1131-1140, 2016.

VIELHAUER, K.; DENICH, M.; KATO, O.R.; DO SOCORRO KATO, M.; DE ABREU SÁ, T.D.; VLEK, P.L.G. Viability of Mechanized Chop-and-Mulch as Soil Improving Alternative to Manual Slash-and-Burn Land Preparation, Rural Poverty Reduction through Research for Development. Deutscher tropentag, Berlin. 2004.

VAN VLIET, N.; MERTZ, O.; HEINIMANN, A.; LANGANKE, T.; SCHMOOK, B.; ADAMS, C.; SCHMIDT-VOGT, D.; MESSERLI, P.; CASTELLA, J.; JØRGENSEN, L.; BIRCH-THOMSEN, T.; BECH-BRUUN, T.; ICKOWITZ, A.; CHI VU, K.; YASUYUKI, K.; PASCUAL, U.; LEISZ, S.; HETT, C.; PADOCH, C.; DRESSLER, W.; ZIEGLER, A.; FOX, F. Trends, drivers and impacts of changes in swidden cultivation in tropical forest-agriculture frontiers: A global assessment. **Global Environmental Change**, v. 22, p. 418–429, 2012.

VASCONCELLOS, R.L.F.; SEGAT, J.C.; BONFIM, J.A.; BARETTA, D.; CARDOSO, E.J.B.N. Soil macrofauna as an indicator of soil quality in an undisturbed riparian forest and recovering sites of different ages. **European Journal of Soil Biology**, v. 58, p. 105–112. 2013.

VERMA, S.; JAYAKUMAR, S. Impact of forest fire on physical, chemical and biological properties of soil: A review. Proceedings of the International Academy of Ecology and Environmental Sciences. v, 2 n. 3, p.168-176, 2012.

WELCHEZ, L.A.; AYARZA, M.; AMEZQUITA, E.; BARRIOS, E.; RONDON, M.; CASTRO, A.; RIVERA, M.; PAVON, J.; FERREIRA, O.; VALLADARES, D.; SANCHEZ, N.; RAO, I.M. No-burn agricultural zones in Honduran hillsides: better harvests, air quality, and water availability by way of improved land management. In: Sustainable Land Management Sourcebook. The World Bank, Agriculture and Rural Development, Washington, DC, p. 78–82, 2008.

CAPÍTULO II

Artigo escrito de acordo com as normas da revista *Agriculture, ecosystems and environment*

Quesungual Agroforestry System conserves soil macrofauna density and richness of the Amazon forest

Jesús Burgos-Guerrero¹, Luis Hernández-García¹, Ernesto Gomez¹, Stefania Triana¹, Julio Cesar Bravo¹, Danielle Celentano¹, Guillaume Xavier Rousseau^{1,2 *}

¹ Agroecology Graduate Program (PPGAgroecologia), State University of Maranhão (UEMA), Campus Universitário Paulo VI, s/n, Tirirical, 65054-970, São Luís, MA, Brazil.

² Biodiversity Research Program (PPBIO) Amazônia Oriental, Ministry of Science, Technology, Innovation and Communication (MCTI), Brazil.

* correspondent author: guilirou@yahoo.ca

Abstract: The Quesungual agroforestry system (QAS) from Central America may be a suitable alternative for small-scale family farming in eastern Amazon, where the intensification of slash-and-burn agriculture threatens environment and livelihood. This is the first experiment to test this system in the Amazon region. To evaluate its impact on soil quality we assessed soil macrofauna abundance and diversity, soil physical and chemical attributes in QAS, slash-and-burn agriculture (SB) and secondary forests (SF) from different ages of fallow in comparison with mature forests (MF) in Alcantara County, Maranhao state. In total, 32 plots (40 X 50 m) were evaluated; data analysis included mixed models, clustering and RDA. Macrofauna community was a very sensitive indicator of land use systems, especially predators, saprophagous organisms and litter transformers. Results indicate that QAS and MF conserve similar density and richness of macrofauna, while SB had a strong negative impact on it. The Quesungual agroforestry system maximizes soil organic carbon accumulation and represents a viable alternative to promote both soil biodiversity conservation and food security in eastern Amazon.

Key words: Macrofauna, secondary forest, richness, density, Amazon

Introduction

Slash-and-burn agriculture or shifting cultivation has been practiced for thousands years in the tropics (Pedroso et al., 2008; Barni et al., 2012). It is considered a sustainable agricultural method in conditions of low population density, high land availability and long fallow periods (Pedroso et al., 2008; Padoch and Pinedo-Vasquez, 2010; Lawrence et al., 2010). However, the increase of population and its demand for agricultural lands are leading agriculture into a process of intensification (Metzger, 2003; Mertz et al., 2009; van Vliet et al., 2012), causing the reduction of fallow periods and making those systems increasingly less productive and sustainable (Ayarza et al., 2010). Indeed slash-and-burn agriculture is often associated with deforestation, global warming, soil degradation and biodiversity loss (Pedroso et al., 2008; Coe et al., 2013). For these reasons, academic institutions, governments and non-governmental organizations are seeking for alternatives to replace its use (IFA, 2001; Coe et al., 2013).

Proposed alternative practices have included alley cropping (Moura et al., 2009), mulching (Vielhauer et al., 2004) no-tillage (Baretta et al., 2014) and agroforestry systems (Pinho et al., 2012). The Quesungual agroforestry system is based on the sustainable management of vegetation, water and soil resources (CIAT, 2009). The system consists in cutting and pruning the natural vegetation to provide permanent soil cover and permit trees and shrubs to regenerate within the crop field. The natural regeneration protects the soil from erosion, produces biomass for additional pruning and allows a fast recovery of forest coverage when the land is abandoned (CIAT, 2009; Pauli et al., 2011). On average, crop rotations last 7 years, after which the land is abandoned (Pauli et al., 2011). The name Quesungual comes from the Honduran indigenous community, in Central America, where it was first identified (Hellin et al., 1999). Although the system was originally created for hillside areas, it seems to be adaptable to the eastern Amazon region, which also has long periods of drought, low fertility soils, susceptibility to erosion and common use of the slash-and-burn agriculture. However, both the productivity and sustainability of this method must be evaluated in the Amazon region.

Soil macrofauna community is considered as a good indicator of soil quality and

agricultural sustainability (Brussaard et al. 2007). These invertebrates play a key role in the provision of soil ecosystem services as their density and diversity are associated with best soil functioning (Lavelle et al., 2006; Rousseau et al., 2014). Indeed, large taxonomic groups of soil macrofauna (class, order or family) are often used to evaluate soil management strategies and the restoration process (Lobry de Bruyn, 1999; Vasconcellos et al., 2013; Rousseau et al., 2013), given that they are strongly influenced by land use systems and produce fast and reliable results (Lavelle et al., 2006; Marichal et al., 2014; Moura et al., 2014). In Central America, Rousseau et al. (2012) found that the abundance of macrofauna groups was a good indicator of soil quality in Agroforestry systems compared to mature forests.

As far as we know it is the first experiment to test the Quesungual agroforestry system in the Amazon region of Brazil. This study aimed to characterize and compare the density and diversity of the soil macrofauna groups in the Quesungual agroforestry system, slash-and-burn agriculture, secondary forest and mature forest in Maranhão, eastern Amazon. Our hypothesis is that the Quesungual agroforestry system can host the same density and diversity of soil invertebrates as that of the conserved forest remnants.

Materials and methods

Study area

The study was conducted in five villages in the county of Alcântara, northern Maranhão state, Brazil (02°19'17" - 02°24'05" S and 44°25'42"- 44°28'49" W) (Figure 1). This region is part of the Eastern Amazon biome (IBGE, 2002); the climate is tropical, according to the classification system of Köppen (Alvares et al., 2013), with two well-defined seasons: a rainy season from January to June and a dry season from July to December. The average annual precipitation varies between 1,000 and 1,800 mm, with an average temperature of 25 °C (Brito and Rego, 2001). The soil is classified as Plinthosol (Anjos et al., 1995) characterized by low fertility and poor cohesion (Moura et al., 2014). The landscape is dominated by young secondary forests, some old forest fragments (Zelarayán et al., 2015) and areas with slash-and-burn agriculture. The population is very poor and relies on the slash-and-burn agriculture for subsistence (Celentano et al., 2014).




Figure 1. Location of study areas in Alcântara County, Maranhão state, Brazil. Land uses: SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forest, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.

2.2. Experimental design

The experimental design was comprised of eight land uses, each one with four replicates: MF - mature forest (> 120 yr-old, with selective logging); YSF - early-succession secondary forests (3 to 6 yr-old); MSF - intermediate-succession secondary forests (7 to 12 yr-old); OSF - advanced-succession secondary forests (30 to 40 yr-old); SBY - slash-and-burn agriculture from early-succession secondary forests; SBM - slash-and-burn agriculture from intermediate-succession secondary forests; QASY - Quesungual agroforestry system from early-succession secondary forests; QASM - Quesungual agroforestry system from intermediate-succession secondary forests. Forest ages (time since last slash-and-burn) were reported by each landowner (Table 1).

Table 1. Description of plots and land uses in Alcântara County, Maranhão state, Brazil.

Land use	Abbreviation	Sampling year	Village	Coordinates		Succession age (years)
				South	West	
Slash-and-burn agriculture from early-succession secondary forests	SBY	2015	Cajueiro	S 02°22'24"	W 44°27'45"	4
	SBY	2015	Espera	S 02°22'13"	W 44°26'42"	3
	SBY	2016	Espera	S 02°22'08"	W 44°26'44"	4
	SBY	2016	Espera	S 02°22'04"	W 44°26'19"	5
Slash-and-burn agriculture from intermediate-succession secondary forests	SBM	2015	Só assim	S 02°21'26"	W 44°28'40"	8
	SBM	2015	Espera	S 02°21'55"	W 44°27'14"	7
	SBM	2016	Só assim	S 02°21'28"	W 44°28'49"	7
	SBM	2016	Espera	S 02°21'53"	W 44°27'15"	7
Quesungual agroforestry system from early-succession secondary forests	QASY	2015	Cajueiro	S 02°22'24"	W 44°27'42"	4
	QASY	2015	Espera	S 02°22'09"	W 44°26'42"	3
	QASY	2016	Espera	S 02°22'09"	W 44°26'42"	3
	QASY	2016	Espera	S 02°22'04"	W 44°26'18"	3
Quesungual agroforestry system from intermediate-succession secondary forests	QASM	2015	Só Assim	S 02°21'27"	W 44°28'40"	8
	QASM	2015	Espera	S 02°21'54"	W 44°27'14"	7
	QASM	2016	Espera	S 02°21'54"	W 44°27'13"	7
	QASM	2016	Só Assim	S 02°21'27"	W 44°28'40"	7
Early-succession secondary forests	YSF	2015	Espera	S 02°22'13"	W 44°26'40"	5
	YSF	2015	Cajueiro	S 02°22'26"	W 44°27'43"	4
	YSF	2016	Espera	S 02°22'02"	W 44°26'23"	4
	YSF	2016	Espera	S 02°22'03"	W 44°26'42"	5
Intermediate-succession secondary forests	MSF	2015	Espera	S 02°21'52"	W 44°27'14"	7
	MSF	2015	Só Assim	S 02°21'29"	W 44°28'39"	8
	MSF	2016	Só Assim	S 02°21'21"	W 44°28'44"	10
	MSF	2016	Espera	S 02°21'51"	W 44°27'16"	10
Advanced-succession secondary forests	OSF	2015	Espera	S 02°21'56"	W 44°27'10"	>30
	OSF	2015	Maruda	S 02°19'17"	W 44°28'28"	>30
	OSF	2016	Pepital	S 02°21'02"	W 44°25'48"	>30
	OSF	2016	Pepital	S 02°20'57"	W 44°25'42"	>30
Mature forest.	MF	2015	Espera	S 02°21'58"	W 44°27'16"	>120
	MF	2015	Cajueiro	S 02°24'02"	W 44°27'40"	>120
	MF	2016	Espera	S 02°21'57"	W 44°27'16"	>120
	MF	2016	Cajueiro	S 02°24'05"	W 44°27'33"	>120

The Quesungual agroforestry and slash-and-burn systems were implemented in 40 x 50 m plots (0.2 ha) in October 2014 and 2015. To prepare the slash-and-burn plots, all vegetation inside the plots were cut, evenly distributed and burned (after two months when it was completely dry). Cassava (*Manihot esculenta Crantz*) and maize (*Zea mays*) were planted at the beginning of the rainy season (March 2015 and February 2016). To prepare the Quesungual plots, trees were pruned to 1.5m of height and those of interest (fruit and timber) were left standing, then the material (leaves and branches) were placed as mulching throughout the plot (FAO, 2005). In March 2015 and February 2016, we planted cassava, maize, pigeon peas (*Cajanus cajan*) and jack beans (*Canavalia ensiformis*) as green manure (Figure 2). Before the sowing the plots were manured with limestone (2 Mg/ha) and rock phosphate (0.24 Mg/ha).




Figure 2. The eight land-use evaluated in Alcântara County, Maranhão state, Brazil. Land uses: SBY = slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.

Soil macrofauna sampling

Sampling was carried out in May 2015 and June 2016 (end of the rainy season). Macrofauna were collected according to the modified Tropical Soil Biology and Fertility method (TSBF, Anderson and Ingram, 1993). Five soil monoliths of 0.25 x 0.25 m and 0.10 m depth, were extracted in a cruciform shape from each plot, with one central monolith and one 20 m apart in each direction (North, South, East and West). The macrofauna (soil invertebrates with 2 mm < body diameter < 20 mm) were

hand-sorted from litter and soil separately, preserved in 99% alcohol and identified as to large taxonomic (class, order or family) and functional groups (according to the dietary or behavioral habits) (Brawn, 2001; Swift et al., 2010). The specimens captured were deposited in the Soil Invertebrates Collection of the Maranhão State University (UEMA).

Soil analysis

Five soil samples were sampled in each plot (one next to each monolith) from 0 to 10 cm depth, with volumetric rings of 282.6 cm³ volume. The soil was weighed wet and divided into sub-samples according to the type of analysis. For determination of moisture content and bulk density, a 40 g soil sub-sample was oven-dried at constant temperature (105 °C for 48 hours) and the moisture percentage was calculated with $((ww-dw) / dw) \times 100$ and the bulk density with dw/v where ww is the wet weight of the sample, dw its dry weight and v the ring volume. For the soil particle-size and chemical analysis, the soil was air-dried and passed through a 2 mm sieve. Textural composition was determined according to the pipette method (Embrapa, 1997). For chemical analysis, we followed the procedures of the Agronomic Institute of Campinas (van Raij et al., 2001) in order to determine the pH (0.01 M suspension in CaCl₂), total organic carbon content (by combustion), P, exchangeable K, Ca, Mg, Na (Resin) and H+Al (SMP method). We also calculated the cation exchange capacity as K+Ca+Mg+(H+Al) and sum of bases as K+Ca+Mg.

Statistical analysis

We calculated the total density of soil macrofauna as the sum of all individuals across taxa per square meter (ind/m²) and assessed diversity through plot richness and mean taxa richness (number of taxa per sample) and the Shannon-Weaver and Pielou indices (adapted from Bandeira et al., 2013). To determine the effect of the land use system on the macrofauna characteristics (density and diversity) and soil properties, we performed generalized linear mixed models (GLMM) by maximum likelihood with negative binomial distribution for all taxa except Auchenorrhyncha and Dermaptera, for which the Poisson distribution was used. In the GLMM, land uses served as the fixed factor whereas the sampling year and sampling location were the random factors. For the choice of most parsimonious models, we used the lowest values of the Akaike information criterion (Di Rienzo et al., 2011). Mean comparison

was performed according to the Tukey test, with significance level of $p \leq 0.05$.

The degree of similarity between the land uses was determined by the Bray-Curtis index with graphic representation as dendrogram. This index of association expresses the similarity between groups, sites or environments based on abundance of species or taxa (Legendre and Legendre, 1998).

The relationship between the soil macrofauna and the soil physical and chemical attributes was studied through Canonical Redundancy Analysis (RDA). This method represents the relationship between biotic and abiotic variables, where X is the matrix of explanatory variables (soil attributes) and Y is the matrix of response variables (soil macrofauna) (Legendre and Legendre, 1998). We used the Forward Selection to select the soil attributes that best explain the macrofauna composition and test their significance. The soil macrofauna density and physico-chemical data were transformed by $\ln(x + 1)$ before performing RDA.

The analyses of diversity and association index were made with the statistical programs InfoStat student version 2016 (Di Rienzo et al., 2011). The mixed models and RDA analyses were performed with the software R (R Development Core Team 2009) using the packages lme4 (GLMM) (Bates et al., 2012) and vegan (RDA and collector curve; Oksanen et al., 2008).

Results

A total of 6,121 individuals were counted and classified into 24 taxonomic groups, of which the most abundant were Isoptera (33.1%), Formicidae (28.8%), Oligochaeta (16.9%), Coleoptera (4.6%), Chilopoda (3.7%), Araneae (3.6%), Diplura (1.4%), Gastropoda (1.2%), and Pseudoscorpionida (1.1%). The other groups represented less than 1% of total abundance (Table A1).

Mature Forest (MF) was the only land use with the presence of all 24 taxa found in the study. According to the collector curve, most of land uses were close to saturation, except in slash-and-burn agriculture from early-succession secondary forests (SBY) that showed an ascending pattern even at the 20th sample (Figure A1).

Soil macrofauna density and diversity

The total density and richness were highest in MF and Quesungual agroforestry system from intermediate-succession secondary forests (QASM), differing significantly from SBY (Table 2). On average, the MF had among 15 to 19 taxa per plot, QASM 14 to 18 and SBY 4 to 9. Other land uses had between 9 and 16 taxa per plot and were ranked as follows: MF > QASM > OSF > MSF > QASY > YSF > SBM > SBY. Although the Shannon-Weaver and Pielou equitability index also followed this trend, they did not differ statistically ($p \leq 0.81$ and $p \leq 0.96$). According to the mixed models, neither the year nor the location influenced the total density or taxa richness.

Comparing the agricultural areas, we found that age of the secondary forest slashed for the implementation of agricultural systems with or without employing fire showed significant influence on the density ($F = 19.5$; $p \leq 0.0001$) and richness ($F = 20.4$; $p \leq 0.0167$) of soil macrofauna.

Table 2. Soil macrofauna community-structure indices (mean and standard errors) in different land uses within Alcântara County, Maranhão state, Brazil.

Community Structure	Land Uses*								F	p
	SBY	SBM	QASY	QASM	YSF	MSF	OSF	MF		
Total Density	246.4 ±91.6 b	436.0 ±91.2 ab	454.4 ±31.0 ab	920.8 ±149.3 a	486.4 ±259.4 ab	529.6 ±150.3 ab	824.8 ±494.3 a	1021.6 ±59.43 a	19.5	≤0.0068
Mean Richness	3.05 ±0.5 b	4.9 ±0.1 ab	6.3 ±0.1 ab	8.2 ±1.2 a	5.0 ±1.5 ab	7.1 ±0.3 ab	6.5 ±0.4 ab	9.8 ±1.1 a	20.4	≤0.0043
Shannon index	0.62 ±0.1	1.12 ±0.02	1.36 ±0.03	1.31 ±0.2	1.1 ±0.04	1.47 ±0.07	1.34 ±0.1	1.64 ±0.1	-	≤0.8146
Pielou index	0.52 ±0.08	0.75 ±0.01	0.74 ±0.01	0.67 ±0.02	0.72 ±0.05	0.77 ±0.03	0.75 ±0.09	0.74 ±0.01	-	≤0.9695

Means with different letter indicate significant differences according to the Tukey test ($p \leq 0.05$). Land uses: SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.

Soil macrofauna composition

Twelve taxa were found in all land use systems (Isoptera, Formicidae, Oligochaeta, Coleoptera, Chilopoda, Araneae, Gastropoda, Pseudoscorpionida, Heteroptera, Diptera, Orthoptera and Lepidoptera), whereas two taxa (Auchenorrhyncha and Scorpionida) were exclusive in MF.

Eight taxa had densities that differed significantly between the land uses (Table A1). Formicidae density was higher in QASM without significant difference versus the majority of land uses except SBY. Oligochaeta was considerably more abundant in MF, the fallows and the Quesungual, while areas with slash-and-burn had the lowest abundances. Predators such as Araneae, Chilopoda and Pseudoscorpionida had the highest density in the MF and QASM. Diplopoda showed maximum density in MF followed by SBM; Gastropoda was more abundant in QASM than in others land uses. Ixodida had a high density in OSF and was not present in SBY or QASY.

The dendrogram based on Bray-Curtis index separated land uses in three groups (distance=1.05): (1) SBY; (2) MF and QASM; and (3) other land uses. The cluster analyses confirmed that QASM was very similar to the most conserved forests and that the SBY has a severely depleted macrofauna community. All other land uses maintained an intermediate status (Figure 3).




Figure 3. Density of soil macrofauna (ind/m²) by taxa and dendrogram of Bray-Curtis index in Alcântara County, Maranhão state, Brazil. Land uses: SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.

Soil attributes and macrofauna

Physical attributes of soil did not statistically differ among land uses. According to the chemical soil attributes, QASM presented the best agronomic quality. Organic Carbon contain ($F = 18.1$; $p \leq 0.0111$), Phosphorous ($F = 18.1$; $p \leq 0.0112$), Magnesium ($F = 22.5$; $p \leq 0.0020$), Cation Exchange Capacity ($F = 26.2$; $p \leq 0.0004$), Potential Acidity ($F = 58.8$; $p \leq 0.0443$) and Sum of Bases ($F = 14.4$; $p \leq 0.0001$) were significantly different between the land uses (Table 3).

Table 3. Physical and chemical attributes of soils (average and standard errors) in different land use in Alcântara County, Maranhão state, Brazil.

Soil attribute	United	SBY*	SBM*	QASY*	QASM*	YSF*	MSF*	OSF*	MF*	F	P value
Chemical attributes											
Organic Carbon	Mg/ha	12.70 ± 0.001 b	15.93 ± 0.002 ab	12.76 ± 0.001 b	19.45 ± 0.003 a	17.38 ± 0.002 ab	18.87 ± 0.003 ab	16.37 ± 0.002 ab	17.27 ± 0.001 ab	18.1	≤ 0.0111
pH	CaCl ₂	4.7 ± 0.21	4.7 ± 0.27	5.1 ± 0.2	5.2 ± 0.2	4.4 ± 0.18	4.2 ± 0.2	4.1 ± 0.2	4.1 ± 0.2	-	0.9913
Phosphorous	Mg/ha	0.02 ± 0.004 b	0.04 ± 0.010 ab	0.04 ± 0.025 ab	0.08 ± 0.046 a	0.02 ± 0.003 b	0.03 ± 0.006 ab	0.02 ± 0.001 ab	0.03 ± 0.006 ab	18.1	≤ 0.0112
Potassium	Mg/ha	0.06 ± 0.004	0.08 ± 0.011	0.06 ± 0.011	0.07 ± 0.010	0.07 ± 0.018	0.07 ± 0.013	0.06 ± 0.008	0.05 ± 0.003	-	0.1962
Calcium	Mg/ha	0.28 ± 0.062	0.41 ± 0.094	0.37 ± 0.062	0.52 ± 0.126	0.31 ± 0.073	0.34 ± 0.136	0.17 ± 0.066	0.31 ± 0.125	-	0.1903
Magnesium	Mg/ha	0.09 ± 0.022 b	0.17 ± 0.043 ab	0.13 ± 0.035 ab	0.21 ± 0.054 a	0.08 ± 0.012 b	0.13 ± 0.032 ab	0.09 ± 0.016 ab	0.09 ± 0.021 ab	22.5	≤ 0.0020
Sodium	Mg/ha	0.05 ± 0.0002	0.09 ± 0.0006	0.07 ± 0.0006	0.08 ± 0.0007	0.07 ± 0.0009	0.08 ± 0.0006	0.08 ± 0.0005	0.04 ± 0.0002	-	0.5837
Cation Exchange Capacity	mmol _c /dm ³	50.59 ± 4.1 c	77.83 ± 12.3 c	58.6 ± 2.3 bc	79.8 ± 12.6 ab	78.6 ± 6.1 ab	90.3 ± 10.1 a	82.3 ± 9.3 ab	83.1 ± 4.1 ab	26.2	≤ 0.0004
Potential Acidity	mmol _c /dm ³	28.05 ± 3.0 b	41.45 ± 11.4 ab	27.9 ± 3.9 b	35.1 ± 7.2 b	54.4 ± 10.0 a	61.1 ± 4.6 a	64.5 ± 9.6 ab	58.6 ± 8.6 a	58.8	≤ 0.0001
Sum of Bases	mmol _c /dm ³	22.5 ± 4.8 ab	36.3 ± 8.2 ab	30.6 ± 6.2 ab	44.6 ± 10.3 a	24.2 ± 4.3 ab	29.1 ± 9.3 ab	17.7 ± 4.6 b	24.5 ± 7.9 ab	14.4	≤ 0.0443
Physical attributes											
Soil moisture	%	8.0 ± 1.2	12.3 ± 2.3	11.2 ± 0.8	11.9 ± 2.5	11.7 ± 2.2	11.9 ± 1.9	10.8 ± 1.5	9.6 ± 2.2	-	0.4283
Bulk Density	g/cm ³	1.40 ± 0.05	1.3 ± 0.04	1.4 ± 0.07	1.3 ± 0.1	1.3 ± 0.02	1.2 ± 0.03	1.2 ± 0.09	1.2 ± 0.05	-	1.0000
Sand	%	75.7 ± 3.4	80.6 ± 2.1	80.2 ± 0.7	77.2 ± 0.7	76.6 ± 1.5	77.4 ± 1.6	80.7 ± 1.6	79.3 ± 1.1	-	0.9635
Silt	%	6.1 ± 0.4	5.1 ± 0.5	7.3 ± 1.6	5.5 ± 0.3	5.6 ± 0.5	5.5 ± 0.3	5.4 ± 0.4	5.1 ± 0.6	-	0.9164
Clay	%	18.2 ± 3.0	14.3 ± 1.7	14.7 ± 0.9	17.3 ± 0.4	17.3 ± 1.3	17.1 ± 1.4	14.5 ± 1.5	15.6 ± 0.7	-	0.7618

Means with different letter indicate significant differences according with Tukey test ($p \leq 0.05$).

* SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.

According to the Forward Selection procedure, only four of the fifteen physico-chemical soil attributes analyzed (Table 3) were significantly correlated with the total density of soil macrofauna (bulk density, $p \leq 0.04$; clay content, $p \leq 0.01$; soil moisture,

$p \leq 0.01$; sand content, $p \leq 0.005$) and were included in the RDA (Figure 4). This soil attributes accounted for 25% of the total variation of soil macrofauna density. Axis 1 accounted for 13.5% of the total variation of soil macrofauna data while Axis 2 explained 6.8%. Soil moisture was the attribute most important for contributing to the first component, while the textural attribute of soil was the second most important variable for explaining the total variance of data, as shown by the second component.




Fig. 4. Canonical redundancy analysis (RDA) with soil macrofauna density, soil physico-chemical and sampling sites in different land-uses in Alcântara County, Maranhão state, Brazil. Land uses: SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest. Taxa: Ara= Araneae, Auch= Auchenorrhyncha, Bla= Blattaria, Col= Coleoptera, Chi= Chilopoda, Der= Dermaptera, Dip= Diptera, Diplo= Dilopoda, Diplu= Diplura, Emb= Embioptera, Fer= Formicidae, Gas= Gastropoda, Lep= Lepidoptera, Het= Heteroptera, Isop= Isoptera, Isopo= Isopoda, Ixo= Ixodida, Neu= Neuroptera, Oli= Oligochaeta, Opi= Opilionida, Other Hym= Others Hymenoptera, Pse= Pseudoscorpionida, Scor= Scorpionida.

Discussion

The development of agro-ecosystems that sustain productivity while promoting biodiversity is a challenge for improving rural livelihoods and achieving conservation

goals (Velasquez et al., 2012). According to Brussaard et al. (2007), the conservation of soil macrofauna constitutes a significant aspect of agricultural sustainability. Indeed, soil macrofauna is more abundant and diverse in land use systems that conserve a structure similar to the original forest (Barros et al, 2002), while land use intensification for agriculture decreases soil macrofauna density and diversity (Barros et al., 2002; Mathieu et al., 2005; Bartz et al., 2014). Soil macrofauna abundance and richness found in our study area were consistent with other studies in the eastern Amazon (Rousseau et al., 2014; Triana et al., 2015).

Slash-and-burn agriculture had a strong negative impact on the richness and density of soil macrofauna (Rossi et al., 2010). The increase of soil surface temperature up to 400°C, the litter layer removal and the soil exposure to direct solar radiation (Matlack, 1993; Gimeno-García et al., 2004), lead to the loss of food sources, ecological niches and high mortality of soil organisms (Rossi et al., 2010). Instead, the Quesungual agroforestry system positively affected the richness and density of soil macrofauna, with values similar to the mature forest. Our results also confirm the ecological importance of fallow period as it influences the quantity and quality of resource available to the soil and the organisms that depend on it (Styger et al., 2007).

Quesungual is characterized by minimum soil disturbance, the continuous accumulation of litter and the presence of trees and shrubs in natural regeneration together with crop field (Pauli et al., 2011; Rousseau et al., 2013), conditions that favor the soil invertebrates (Menezes et al., 2009; Pauli et al., 2011). Although accelerating decomposition processes of tropical soils tends to diminish differences between soil management practices (Moura et al., 2014), the Quesungual system showed better soil attributes as it allowed greater deposition of biomass in the soil, similar to mature forest. The improved Organic Carbon of the soil is very important to conserve soil organisms, guarantee soil productivity and deliver soil ecosystem services (Moura et al., 2014; Celentano et al., 2016).

Although the difference in soil moisture and soil bulk density were not significant, their effect is narrowly associated with the best development of soil macrofauna communities. Land use systems with capacity to maintain soil moisture and no

increase in soil bulk density are very important to maintain soil organism populations (Celentano et al., 2016).

The macrofauna taxa that showed statistical differences between land use systems are those related to a stable environment, with high availability of food resources and a broad supply of ecological niches (Tews et al., 2004; Sayer et al., 2010). Predators (Araneae, Chilopoda and Pseudoscorpionida) were highly sensitive to changes in land use. Indeed, these groups are often related to conserved ecosystems with higher content of organic matter (Menezes et al., 2009; Vasconcellos et al., 2013; Moura et al., 2014), indicating the normal flow of energy and regular functioning of the trophic chain within the communities (Neutel et al., 2002; Moura et al., 2014).

The litter transformers and saprophages (organisms that feed on plant material in decomposition) are very important in the soil functions, especially nutrient cycling (Denich et al., 2004; Lavelle et al., 2006; Moura et al., 2014). The litter transformers such as Gastropoda are strongly associated with the quantity and quality of plant material deposited on the soil, preferring environments with greater quantity and diversity of plants (Rueda-Ramírez and Varela, 2016). While the saprophagous organisms, particularly diplopods are among the easiest to lose after a burning process and one of the most difficult to recover in restoration processes (Korobushkin et al., 2017).

High density Oligochaeta are commonly found in environments with largest soil moisture, less disturbed soil and the presence of organic soil cover. These organisms improve soil processes such as nutrient cycling, aggregation, water infiltration and aeration (Bartz et al., 2014).

The Quesungual system conserves macrofauna biodiversity and promotes soil fertility. According to Pauli et al. (2011) results in Central America, this system can also extend time of crop production (between 7 to 12 years), thus augmenting fallow periods and diminishing pressure throughout the forest. The Quesungual system has the potential to replace slash-and-burn agriculture in the eastern Amazon, with many ecological benefits. The widespread adoption of the Quesungual in Central America (Pauli et al., 2011; Rousseau et al., 2013) suggests that the smallholders found it well

adapted to their needs. Preliminary results of cassava harvesting in Alcântara showed that the Quesungual system had similar yields to the slash-and-burn plots (data not presented) and that eight families involved in the project are interested in maintaining Quesungual testing plots independently. This social interest in agroecological transition must be confirmed, along with the ability of the Quesungual areas to restore soils and sustain demanding crops such as maize that assure food security in rural communities of the eastern Amazon.

Conclusions

The preliminary results of the use of the Quesungual agroforestry system in eastern Amazon suggest that this system represents a suitable alternative to replace slash-and-burn agriculture in small-scale family farming, with notable environmental gains. Soil macrofauna community was a very sensitive indicator of land use systems and it confirmed the ecological importance of fallow period as it influences the quantity and quality of resource available to the soil and the organisms that depend on it. Quesungual agroforestry system conserved similar density and richness of macrofauna in relation to mature forests allowing the normal flow of energy and regular functioning of the trophic chain within the communities, while slash-and-burn agriculture had strong negative impact on soil macrofauna community and soil quality as a consequence. The Quesungual agroforestry system maximized soil organic carbon accumulation and soil protection, suggesting a viable alternative to promote both soil biodiversity conservation and food security in eastern Amazon.

Acknowledgments

We thank the Maranhão State Foundation for Research Development (FAPEMA), the Brazilian Council for Higher Education (CAPES) and UEMA for the financial support. Thanks also to Alcântara communities for their collaboration with soil macrofauna sampling.

References

- Alvares, C. A., Stape, J. L., Sentelhas, P. C., Moraes-Gonçalves, J. L., Sparovek, G., 2013. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, 22, 711-728.
- Anderson, J.M., J.S.I. Ingram., 1993. *Tropical Soil Biology and Fertility: a handbook of methods*, 2a ed. Eds. Anderson and Ingram. CAB International, Wallingford, UK. pp. 89-112.
- Anjos, LHC., Franzmeier DP., Schulze DG., 1995. Formation of soils with plinthite on a toposequence in Maranhão state, Brazil. *Geoderma*. 64, 257-279.
- Ayarza, M., Huber-Sannwald, E., Herrick, J., Reynolds, J., Garcia-Barrios, L., Welchez, L., Lentos, P., Pavo, J., Alvarado, A., Pinedo, M., Baquera, N., Zelaya, S., Morales, J., Pineda, R., Amezcuita, E and Trejo, M., 2010. Changing human–ecological relationships and drivers using the Quesungual agroforestry system in western Honduras. *Renew. Agricult. Food. Syst.* 1-9.
- Bandeira, B., Jamet, J., Jamet, D., Ginoux, J., 2013. Mathematical convergences of biodiversity indices. *Ecol indicat.* 29, 522-528.
- Baretta, D., Bartz, M., Fachini, I., Anselmi, R., Zortéa, T., Baretta, C., 2014. Soil fauna and its relation with environmental variables in soil management systems. *Rev. Ciênc Agron.* 45 (5) 871-879.
- Barni, P. E., Fearnside, P. M., Graça, P. M. L. A., 2012. Desmatamento no sul do Estado de Roraima: padrões de distribuição em função de Projetos de Assentamento do INCRA e da distância das principais rodovias (BR-174 e BR-210). *Acta Amazônica*, 42 (2), 195-204.
- Barros, E., Pashanasi B., Constantino, R., Lavelle, P. 2002. Effects of land-use system on the soil macrofauna in western Brazilian Amazonia. *Biol. Fert. Soils*, 35, 338-347.
- Bartz, C., Brawn, G.G., Gonçalves da Rosa, M., Folho, O., James, S., Decaëns, T., Baretta, D. 2014. Earthworm richness in land use systems in Santa Catarina, Brazil. *Appl. soil ecol*, 83, 59-70.
- Bates, D., Maechler, M., Bolker, B, Walker, S. 2012. lme4, Linear mixed-effects models using Eigen and S4. R package version a.a-7.
- Brito, C. M. S., Rego, M. M. C., 2001. Community of male Euglossini bees (Hymenoptera: Apidae) in a secondary forest, Alcântara, MA, Brazil. *Brazilian J. Biol*, 61(4), 631-638.
- Brown, G. G., fragoso, C., Barois, I., Rojas, P., Patrón, J. C., Bueno, J., Moreno, A. G., Lavelle, P., Ordaz, V., Rodríguez, C., 2001. Diversidad y rol funcional de la macrofauna edáfica en los ecosistemas tropicales mexicanos. *Acta. Zool. Mexicana Esp.* 1, 79-110.

- Brussaard, L., de Ruiter, P.C., Brown, G.G., 2007. Soil biodiversity for agricultural sustainability. *Agric. Ecosyst. Environ.* 121, 233–244.
- Celentano, D., Rousseau, G.X., Engel, V.L., Façanha, C.L., De Oliveira, E.M., De Moura, E.G., 2014. Perceptions of environmental change and use of traditional knowledge to plan riparian forest restoration with relocated communities in Alcantara, Eastern Amazon. *J. Ethn. Ethnomedic.* 10(11), 4-15.
- Celentano, D., Rousseau, G.X., Engel, V.L., Zelarayán M., Oliveira, E.C., Araujo, A.C., Moura, E.G., 2016. Degradation of riparian forest affects soil properties and the provision of ecosystem services in eastern amazon, brazil. *Land Degrad Develop.* 12 p.
- CIAT- International Center for Tropical Agriculture, 2009. Quesungual slash and mulch agroforestry system (QSMAS): Improving crop water productivity, food security and resource quality in the subhumid tropics. CPWF Project Report. Cali, Colombia. pp. 1-64.
- Coe, M.; Marthews, T.; Costa, M.; Galbraith, D.; Greenglass, N.; Imbuzeiro, H.; Levine, N.; Malhi, Y.; Moorcroft, P.; Muza, M.; Powell, T.; Saleska, S.; Solorzano, L.; Wang, J., 2013. Deforestation and climate feedbacks threaten the ecological integrity of south –southeastern Amazonia. *Phil. Trans. R. Soc. B.* 368, 1-9.
- Denich, M., Vielhauer, K., de A. Kato, M.S., Block, A., Kato, O.R., de Abreu Sá, T.D., Lücke, W., Vlek, P.L.G., 2004. Mechanized land preparation in forest-based fallow systems: the experience from Eastern Amazonia. *Agroforest. Syst.* 61–62, 91–106.
- Di Rienzo J.A., Casanoves F., Balzarini M.G., Gonzalez L., Tablada M., Robledo C.W., 2011. InfoStat versión 2016. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina.
- EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária., 1997. Manual de métodos de análise de solos. Centro Nacional de Pesquisas de Solos, Rio de Janeiro. pp. 212.
- FAO - Food and Agriculture Organization of the United Nations, 2005. El sistema agroforestal Quesungual: Una opción para el manejo de suelos en zonas secas de ladera. pp. 1-50.
- IFA - Fertilizer Industry Association, 2001. Global estimates of gaseous emissions of NH₃, NO and N₂O from agricultural land. Primera versión. Roma, Italia. pp. 106.
- Gimeno-Garcia, E., Andreu, V., Rubio, J., 2004. Spatial patterns of soil temperatures during experimental fires. *Geoderma* 118, 17–38.
- Hellin, J., Welchez, L.A., Cherrett, I., 1999. The Quesungual System: an indigenous agroforestry system from western Honduras. *Agroforest. Syst.* 46, 229-237.

- IBGE - Instituto Brasileiro de Geografia e Estatística, 2002. Resolução N° 05, 10 de outubro de 2002. Diário oficial da república federativa do Brasil. Poder ejecutivo, Brasília DF. 11 outubro, 2002. Seção 1. pp. 46-65.
- Korobushkin, D.I., Gorbunova, A.Yu., Zaitsev, A.S., Gongalsky, K.B., 2017. Trait-specific response of soil macrofauna to forest burning along a macrogeographic gradient. *Appl. soil ecol.* 112, 97-100.
- Lavelle, P., Decaëns, T., Aubert, M., Barot, S., Blouin, M., Bureau, F., Margarie, P., Mora, P., Rossi, J.P., 2006. Soil invertebrates and ecosystem services. *Eur. J. Soil Biol.* 42, 3-15.
- Lawrence, D., Radel, K. Tully., B. Schmoock., L. Schneider., 2010. Untangling a decline in tropical forest resilience: constraints on the sustainability of shifting cultivation across the globe. *Biotropica.* 42, 21–30.
- Legendre, P., Legendre, L., 1998. Numerical ecology. Second english edition. Amsterdam. Elsevier. pp. 856.
- Lobry de Bruyn, L.A., 1999. Ants as bioindicators of soil function in rural environments, *Agric. Ecosyst. Environ.* 74, 411–425.
- Marichal, R., Grimaldi, M., Feijoo, A., Oszwald, J., Praxedes, C., Ruiz-Cobo, D., Hurtado, M., Desjardins, T., Lopes, M., Gonzaga, Luis., Souza, I., Nascimento, M., Brown, G., Tsélouiko, S., Martins, M., Decaëns, T., Velasquez, E., Lavelle, P., 2014. Soil macroinvertebrate communities and ecosystem services in deforested landscapes of Amazonia. *Appl. Soil Ecol.* 83. 177–185.
- Mathieu, J., Rossi, J.-P., Mora, P., Lavelle, P., Martins, P.F.D.S., Rouland, C., Grimaldi, M., 2005. Recovery of soil macrofauna communities after forest clearance in Eastern Amazonia. *Brazil. Conserv. Biol.* 19, 1598–1605.
- Matlack, G.R., 1993. Microenvironment variation within and among forest edge sites in the eastern United States. *Biol. Conserv.* 66, 185–194.
- Menezes, C., Fernandes, M., Pereira, M., Batista, I., Couto, W., Cunha, L., Pinto, I., 2009. Macrofauna edáfica em estádios sucessionais de floresta estacional semidecidual e pastagem mista em pinheiral (RJ). *Rev. Bras. Ciênc. solo*, 33 (6), 1647-1656.
- Mertz, O., Padoch, C., Fox, J., Cramb, R.A., Leisz, S.J., Lam, N.T., Vien, T.D., 2009. Swidden change in southeast Asia: understanding causes and consequences. *Human. Ecol.* 37, 259–264.
- Metzger, J. P., 2003. Effects of slash and burn fallow periods on landscapes structure. *Environ. Conserv.* 30 (4), 325-333.
- Moura, E.G., Moura. N.G., Aguiar, A.C.F., 2009. Evaluating chemical and physical quality indicators for a structurally fragile tropical soil. *Soil. Use. Manage.* 25, 368–375.

- Moura, E.G., Aguiar, A.C.F., Piedade, A.R., Rousseau, G.X., 2014. Contribution of legume tree residues and macrofauna to the improvement of abiotic soil properties in the easter Amazon. *Appl. Soil. Ecol.* 86, 91-99.
- Neutel, A.M., Heesterbeek J.A. P., De Ruiter, P.C., 2002. Stability in real food webs: weak links in long loops. *Science.* 296, 1120-1123.
- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G.L., Stevens, M.H.A., Wagner, H., 2008. *Vegan: Community Ecology Package.*
- Padoch, C., Pinedo-Vasquez, M., 2010. Saving slash and burn to save biodiversity. *Biotropica.* 42(5), 550-552.
- Pauli, N., Barrios, E., Conacher, A.J., Oberthür, T., 2011. Soil macrofauna in agricultural landscapes dominated by the Quesungual slash-and-mulch agroforestry system, Western Honduras. *Appl. Soil. Ecol.* 47, 119–132.
- Pedroso J., Murrieta, R., Adams, C. A., 2008. Agricultura de corte e queima: um sistema em transformação. *Boletim Museu do Para Emílio Goeldi, Ciências Humanas*, Belém, 3 (2), 153-174.
- Pinho R. C., Miller R. P., Alfaia S. S., 2012. Agroforestry and the improvement of soil fertility: a view from Amazonia. *Appl. Environ. Soil. Sci.*
- R Development Core Team., 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Austria, <http://www.R-project.org>.
- Rossi J.P., Celini, L., Mora, P., Mathieu J., Lapied, E., Nahmani, J., Ponge, J., Lavelle.P., 2010. Decreasing fallow duration in tropical slash-and-burn agriculture alters soil macroinvertebrate diversity: A case study in southern French Guiana. *Agric. Ecosyst. Environ.* 135, 148–154.
- Rousseau G.X., Deheuvels, O., Rodriguez, I., Somarriba, E., 2012. Indicating soil quality in cacao-based agroforestry systems and old-growth forests: The potential of soil macrofauna assemblage. *Ecol. Indicat.* 23, 535–543.
- Rousseau G.X., P.R. dos S. Silva., C.J.R. de Carvalho., 2014. Impacto do histórico de uso na composição da macrofauna do solo em uma cronosequência de capoeiras, florestas e pastos no Centro de Endemismo Belém, Amazônia Oriental. *Acta Amazônica.* 44 (4): 499-512.
- Rousseau, L., Fonte, S., Van der Hoek, R., Lavelle, P., 2013. Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. *Ecol. Indicat.* 27, 71–82.
- Rueda-Ramírez, D., Varela, A., 2016. Distribución espacial, composición y densidad de edafofauna en hojarasca de bosque y cafetal (montenegro, colombia). *Acta biol. Colombiana.* 21(2). 399-412.

- Sayer, E.J., Sutcliffe, L.M.E., Ross, R.I.C., Tanner, E.V.J., 2010. Arthropod Abundance and Diversity in a Lowland Tropical Forest Floor in Panama: The Role of Habitat Space vs. Nutrient Concentrations. *Biotropica*, 42, 194–200.
- Styger, E., Rakotondramasy, H., Pfeffer, M., Fernandes, E., Bates, D., 2007. Influence of slash-and-burn farming practices on fallow succession and land degradation in the rainforest region of Madagascar. *Agric. Ecosyst. Environ.* 119, 257–269.
- Swift, M. J., Bignell, D., Moreira, F. M. de S., Huisling, J., 2010. O inventário da biodiversidade biológica do solo: conceitos e orientações gerais. Manual de biologia dos solos tropicais: amostragem e caracterização da biodiversidade. pp. 23-41.
- Tews, J.; Brose, U.; Grimm, U.; Tielbörger, K.; Wichmann, M.C.; Schawger, M., Jeltsch, F., 2004. Animal species diversity driven by habitat heterogeneity/diversity: The importance of keystone structures. *J. Biogeogr.* 31, 79-92.
- Triana, S. P., Rousseau, G.X., Rocha da Piedade, A., Celentano, D. Zelarayán, M. L. C., Braun, H., 2015. La macrofauna del suelo como indicadora de degradación de bosques ribereños en la amazonia oriental brasileira. *Rev. Facul. Agron.* 114 (1), 49-60.
- van Raij, B., Andrade, J.C., Cantarella, H., Quaggio, J.A., 2001. Análise Química para Avaliação da Fertilidade de solos Tropicais. Campinas, Instituto Agrônomo.
- Van Vliet, N., Mertz, O., Heinemann, A., Langanke, T., Pascual, U., Schmook, B. et al., 2012. Trends, drivers and impacts of changes in swidden cultivation in tropical forest-agriculture frontiers: a global assessment. *Glob. Environ. Change*, 22, 418–429.
- Vasconcellos, R.L.F., Segat, J.C., Bonfim, J.A., Baretta, D., Cardoso, E.J.B.N., 2013. Soil macrofauna as an indicator of soil quality in an undisturbed riparian forest and recovering sites of different ages. *Eur. J. Soil. Biol.* 58, 105–112.
- Vielhauer, K., Denich, M., Kato, O.R., do Socorro Kato, M., De Abreu Sá, T.D., Vlek, P.L.G., 2004. Viability of Mechanized Chop-and-Mulch as Soil Improving Alternative to Manual Slash-and-Burn Land Preparation, Rural Poverty Reduction through Research for Development. Deutscher tropentag, Berlin.
- Velasquez, E., Fonte, S., Barot, S., Grimaldi, M., Desjardins, T., Lavelle, P., 2012. Soil macrofauna-mediated impacts of plant species composition on soil functioning in amazonia pastures. *Appl. Soil. Ecol.* 56, 43-50.
- Zelarayán, M.L.C., Celentano, D., Oliveira, E.C., Triana, S.P., Muchavisoy, K.H., Rousseau, G.X., 2015. Impacto da degradação sobre o estoque total de carbono de florestas ripárias na Amazônia Oriental, Brasil. *Acta Amazônica*, 45 (3), 271-282.

SUPPLEMENTARY MATERIAL

Table A.1. Means of density (Ind/m²) of the soil macrofauna by taxa and functional groups (average and standard errors) under different land uses in Alcântara County, Eastern Amazon.

Taxonomic Group	SBY*	SBM*	QASY*	QASM*	YSF*	MSF*	OSF*	MF*	F	P-value
Soil engineers										
Formicidae	34.4 ±18.5 c	125.6 ±41.3 abc	155.2 ±14.1 abc	407.2 ±50.7 a	188 ±155.0 ab	74.4 ±21.3 bc	124 ±66.8 abc	304.8 ±101.5 ab	30.2	≤0.0001
Isoptera	152 ±82.7	157.6 ±94.1	122.4 ±41.2	135.2 ±77.7	99.2 ±17.8	192.8 ±147.1	502.4 ±438.9	261.6 ±73.8	-	0.2487
Oligochaeta	28 ±23.7 c	56.8 ±25.4 bc	70.4 ±16.6 abc	152 ±47.5 ab	115.2 ±67.3 ab	125.6 ±21.4 ab	91.2 ±44.4 abc	189.6 ±60.8 a	39.4	≤0.0001
Saprophages										
Diptera	0 ±0	1.6 ±0.9	0.8 ±0.8	5.6 ±2.4	6.4 ±2.9	1.6 ±0.9	1.6 ±0.9	2.4 ±1.5	-	0.2788
Diplopoda	0.8 ±0.8 b	11.2 ±4.8 ab	0.8 ±0.8 b	4.0 ±2.4 ab	0 ±0 c	4.8 ±2.0 ab	2.4 ±1.5 ab	16 ±4.5 a	17.6	≤0.0139
Isopoda	0 ±0	2.4 ±1.5	0.8 ±0.8	4.0 ±4.0	0.8 ±0.8	0 ±0	3.2 ±1.3	16.8 ±8.4	-	0.1616
Litter Transformers										
Embioptera	0 ±0	0 ±0	4.0 ±2.0	3.2 ±1.3	1.6 ±0.9	1.6 ±0.9	2.4 ±1.5	2.4 ±1.5	-	0.2522
Gastropoda	0.8 ±0.8 b	4.0 ±4.0 ab	4.0 ±0.8 ab	22.4 ±14.0 a	5.6 ±2.7 ab	12.8 ±8.5 ab	0.8 ±0.8 b	6.4 ±2.6 ab	20.6	≤0.0043
Heteroptera	0.8 ±0.8	1.6 ±0.9	1.6 ±0.9	8.0 ±6.9	0.8 ±0.8	0.8 ±0.8	4.8 ±1.6	4.8 ±2.0	-	0.1461
Auchenorrhyncha	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0.8 ±0.8	-	1.0000
Lepidoptera	0.8 ±0.8	2.4 ±0.8	1.6 ±1.6	4.0 ±2.0	0.8 ±0.8	1.6 ±1.6	1.6 ±0.9	4.0 ±3.0	-	0.7255
Orthóptera	1.6 ±1.6	0.8 ±0.8	3.2 ±1.3	5.6 ±3.5	0.8 ±0.8	1.6 ±1.6	3.2 ±2.2	4.0 ±2.0	-	0.6446
Omnivores										
Blattária	0 ±0	0.8 ±0.8	4.0 ±2.0	14.4 ±9.1	0 ±0	2.4 ±1.5	2.4 ±2.4	5.6 ±2.0	-	0.2528
Coleoptera	10.4 ±5.7	36.0 ±12.8	32.0 ±5.0	28.8 ±8.5	28.0 ±15.5	24.8 ±4.9	20.8 ±4.0	44.8 ±2.2	-	0.1535
Predators										
Araneae	9.6 ±2.9 c	20.8 ±4.2 abc	24.0 ±7.4 abc	34.4 ±7.9 ab	9.6 ±5.3 c	21.6 ±3.7 abc	12.8 ±4.7 bc	44.8 ±12.4 a	36.3	≤0.0001
Chilopoda	3.2 ±2.2 d	6.4 ±1.8 cd	17.6 ±3.8 abc	44.8 ±18.2 ab	12.8 ±7.5 bcd	28.0 ±6.0 ab	15.2 ±1.5 abc	51.2 ±3.9 a	53.4	≤0.0001
Dermáptera	0 ±0	0 ±0	0 ±0	3.2 ±3.2	2.4 ±1.5	0 ±0	0 ±0	1.6 ±0.9	-	0.7009
Diplura	0 ±0	2.4 ±1.5	2.4 ±1.5	14.4 ±9.2	7.2 ±1.5	12.8 ±3.6	17.6 ±5.3	14.4 ±6.7	-	0.2185
Neuroptera	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0.8 ±0.8	0 ±0	2.4 ±1.5	-	0.2216
Opilionida	0 ±0	0.8 ±0.8	0 ±0	5.6 ±4.5	0 ±0	4.8 ±3.0	8.0 ±4.8	24.0 ±20.8	-	0.8380
Pseudoscorpionida	1.6 ±1.6 ab	0.8 ±0.8 b	8.8 ±0.8 ab	14.4 ±5.9 a	3.2 ±2.2 ab	11.2 ±2.7 a	2.4 ±0.8 ab	13.6 ±5.4 a	25.3	≤0.0006
Scorpionida	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0 ±0	0.8 ±0.8	-	1.0000
Others										
Ixórida	0 ±0 c	0.8 ±0.8 b	0 ±0 c	0.8 ±0.8 b	0.8 ±0.8 b	1.6 ±0.9 ab	4.8 ±2.7 a	0.8 ±0.8 b	355.9	≤0.0001
Others Hymenoptera	0.8 ±0.8	0.8 ±0.8	0.8 ±0.8	0.8 ±0.8	1.6 ±0.9	0 ±0	0 ±0	0.8 ±0.8	-	0.9998

Means with different letter indicate significant differences according to the Tukey test ($p \leq 0.05$).

* SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn

agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forests, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.




Fig.A.1. Taxa accumulation curve of soil macrofauna under different land uses in Alcântara County, Eastern Amazon. Where, SBY= slash-and-burn agriculture from early-succession secondary forests, SBM= slash-and-burn agriculture from intermediate-succession secondary forests, QASY= Quesungual agroforestry system from early-succession secondary forests, QASM= Quesungual agroforestry system from intermediate-succession secondary forest, YSF= early-succession secondary forests, MSF= intermediate-succession secondary forests, OSF= advanced-succession secondary forests, MF= Mature forest.